

**Table A4. Cases of HIV infection (not AIDS) for female adults and adolescents, by exposure category and race/ethnicity, reported through December 2002—39 areas with confidential name-based HIV infection reporting**

Exposure category	White, not Hispanic				Black, not Hispanic				Hispanic			
	2002		Cumulative through 2002 <sup>a</sup>		2002		Cumulative through 2002 <sup>a</sup>		2002		Cumulative through 2002 <sup>a</sup>	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Injection drug use	345	18	3,160	26	609	9	5,701	15	281	15	1,132	17
Hemophilia/coagulation disorder	0	0	15	0	5	0	26	0	2	0	5	0
Heterosexual contact:	729	38	5,286	43	2,443	34	15,711	42	707	37	2,845	43
<i>Sex with injection drug user</i>	150	8	1,539	13	326	5	3,024	8	112	6	662	10
<i>Sex with bisexual male</i>	46	2	488	4	99	1	929	2	23	1	101	2
<i>Sex with person with hemophilia</i>	5	0	101	1	9	0	53	0	2	0	7	0
<i>Sex with HIV-infected transfusion recipient</i>	6	0	45	0	17	0	86	0	14	1	33	0
<i>Sex with HIV-infected person, risk not specified</i>	522	27	3,113	26	1,992	28	11,619	31	556	29	2,042	31
Receipt of blood transfusion, blood components, or tissue	5	0	142	1	37	1	297	1	12	1	43	1
Other/risk not reported or identified	826	43	3,586	29	4,040	57	15,884	42	895	47	2,628	40
<b>Total</b>	<b>1,905</b>	<b>100</b>	<b>12,189</b>	<b>100</b>	<b>7,134</b>	<b>100</b>	<b>37,619</b>	<b>100</b>	<b>1,897</b>	<b>100</b>	<b>6,653</b>	<b>100</b>

Exposure category	Asian/Pacific Islander				American Indian/Alaska Native				Totals			
	2002		Cumulative through 2002 <sup>a</sup>		2002		Cumulative through 2002 <sup>a</sup>		2002		Cumulative through 2002 <sup>a</sup>	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Injection drug use	1	1	13	5	17	33	96	33	1,253	11	10,123	18
Hemophilia/coagulation disorder	0	0	0	0	0	0	1	0	7	0	47	0
Heterosexual contact:	25	34	115	44	21	40	122	42	3,925	35	24,136	42
<i>Sex with injection drug user</i>	0	0	10	4	3	6	46	16	591	5	5,289	9
<i>Sex with bisexual male</i>	1	1	5	2	3	6	8	3	172	2	1,536	3
<i>Sex with person with hemophilia</i>	0	0	0	0	0	0	2	1	16	0	164	0
<i>Sex with HIV-infected transfusion recipient</i>	1	1	2	1	0	0	1	0	38	0	167	0
<i>Sex with HIV-infected person, risk not specified</i>	23	32	98	37	15	29	65	22	3,108	28	16,980	30
Receipt of blood transfusion, blood components, or tissue	0	0	6	2	0	0	0	0	54	0	490	1
Other/risk not reported or identified	47	64	130	49	14	27	72	25	5,822	53	22,580	39
<b>Total</b>	<b>73</b>	<b>100</b>	<b>264</b>	<b>100</b>	<b>52</b>	<b>100</b>	<b>291</b>	<b>100</b>	<b>11,061</b>	<b>100</b>	<b>57,376</b>	<b>100</b>

Note. Includes only persons with HIV infection that has not progressed to AIDS.

Since 2002, the following 39 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, American Samoa, Guam, Northern Mariana Islands, and the U.S. Virgin Islands.

Connecticut has confidential name-based HIV infection reporting only for pediatric cases.

Florida (since July 1997) has had confidential name-based HIV infection reporting only for new diagnoses.

Pennsylvania (October 2002) implemented confidential name-based HIV infection reporting only in areas outside the city of Philadelphia.

Texas (February 1994 through December 1998) reported only pediatric HIV infection cases.

<sup>a</sup> Includes persons with a diagnosis of HIV infection (not AIDS), reported from the beginning of the epidemic through 2002. Cumulative total includes 360 females of unknown or multiple race.